

Pinellas County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Pinellas County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	2.8	(1.5 - 4.1)	3.3	(3.0 - 3.5)			+
Smoked cigars on one or more of the past 30 days	2.9	(1.4 - 4.3)	3.9	(3.6 - 4.2)			+
Used smokeless tobacco on one or more of the past 30 days	1.2	(0.3 - 2.1)	2.3	(2.1 - 2.6)		+	
Used any form of tobacco on one or more of the past 30 days	4.9	(3.0 - 6.8)	6.4	(6.0 - 6.8)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	72.6	(68.8 - 76.3)	72.0	(71.2 - 72.8)			+
Exposed to secondhand smoke during the past 7 days	42.6	(38.3 - 46.9)	41.3	(40.4 - 42.2)			+
Smoking is allowed in the home	11.6	(8.9 - 14.4)	9.9	(9.5 - 10.4)			+
Taught about tobacco use in any classes this year	63.9	(59.4 - 68.4)	53.4	(52.0 - 54.7)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	63.6	(59.8 - 67.5)	63.9	(63.1 - 64.6)			+
Served as an active member of SWAT	3.2	(1.8 - 4.7)	4.3	(3.9 - 4.7)			+
<u>Definitely did not think</u> that smokers have more friends	34.5	(30.3 - 38.7)	33.7	(32.9 - 34.5)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	73.8	(70.0 - 77.5)	73.7	(73.0 - 74.4)			+

Pinellas County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Pinellas County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	21.2	(17.9 - 24.6)	20.1	(19.5 - 20.7)			+
Current asthma, among all individuals	14.3	(11.4 - 17.2)	12.0	(11.5 - 12.5)			+
Asthma attack in past year, among individuals with lifetime asthma	18.7	(11.6 - 25.8)	18.1	(16.8 - 19.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	8.1	(5.7 - 10.5)	11.6	(11.0 - 12.1)		+	
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	16.2	(12.8 - 19.6)	16.5	(15.8 - 17.2)			+
Sufficient vigorous activity	71.9	(68.2 - 75.6)	70.1	(69.2 - 70.9)			+
Sufficient moderate activity	29.6	(26.0 - 33.2)	24.8	(24.1 - 25.4)	+		
Exercised to lose weight or to keep from gaining weight during the past 30 days	41.5	(37.0 - 45.9)	44.4	(43.6 - 45.2)			+
Described themselves as slightly or very overweight	29.4	(25.5 - 33.4)	30.2	(29.4 - 30.9)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	7.9	(5.4 - 10.5)	11.4	(10.8 - 12.0)		+	
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	15.0	(12.1 - 18.0)	17.8	(17.2 - 18.5)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	6.2	(4.3 - 8.0)	7.8	(7.3 - 8.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	12.8	(9.7 - 15.9)	12.6	(12.0 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	20.4	(16.9 - 23.8)	19.2	(18.5 - 19.8)			+

Pinellas County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Pinellas County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	13.0	(10.1 - 15.9)	10.1	(9.6 - 10.6)			+
Smoked cigars on one or more of the past 30 days	10.7	(8.3 - 13.1)	11.4	(10.9 - 11.9)			+
Used smokeless tobacco on one or more of the past 30 days	4.1	(2.4 - 5.9)	5.6	(5.2 - 6.0)			+
Used any form of tobacco on one or more of the past 30 days	19.7	(16.2 - 23.2)	17.9	(17.3 - 18.6)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	57.8	(53.1 - 62.5)	58.4	(57.6 - 59.3)			+
Exposed to secondhand smoke during the past 7 days	55.0	(50.2 - 59.9)	47.6	(46.7 - 48.4)	+		
Smoking is allowed in the home	7.8	(5.1 - 10.4)	10.0	(9.5 - 10.5)			+
Taught about tobacco use in any classes this year	40.5	(34.5 - 46.5)	33.4	(32.4 - 34.4)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	52.5	(47.8 - 57.1)	50.4	(49.6 - 51.2)			+
Served as an active member of SWAT	3.5	(1.7 - 5.3)	4.0	(3.7 - 4.3)			+
<u>Definitely did not think</u> that smokers have more friends	32.4	(28.6 - 36.1)	30.5	(29.8 - 31.2)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	70.1	(66.1 - 74.1)	71.7	(71.0 - 72.4)			+

Pinellas County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Pinellas County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	20.9	(17.0 - 24.8)	20.8	(20.2 - 21.5)			+
Current asthma, among all individuals	10.1	(6.9 - 13.2)	10.6	(10.1 - 11.1)			+
Asthma attack in past year, among individuals with lifetime asthma	18.1	(9.6 - 26.6)	16.3	(15.1 - 17.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	6.9	(4.4 - 9.3)	11.1	(10.5 - 11.6)		+	
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	10.3	(7.5 - 13.0)	14.3	(13.7 - 14.9)		+	
Sufficient vigorous activity	62.9	(57.9 - 67.9)	62.7	(62.0 - 63.5)			+
Sufficient moderate activity	29.1	(24.9 - 33.3)	26.1	(25.4 - 26.7)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	40.4	(35.9 - 44.9)	42.3	(41.5 - 43.1)			+
Described themselves as slightly or very overweight	27.1	(23.6 - 30.6)	29.8	(29.0 - 30.5)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	9.6	(6.6 - 12.6)	11.2	(10.5 - 11.8)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	25.5	(21.6 - 29.5)	23.1	(22.4 - 23.8)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	3.9	(1.8 - 6.1)	6.9	(6.4 - 7.3)		+	
Purposely hurt themselves without wanting to die, during the past 12 months	13.8	(10.2 - 17.4)	12.7	(12.1 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	23.7	(19.5 - 28.0)	24.5	(23.8 - 25.2)			+

Pinellas County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Pinellas County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	8.2	(6.2 - 10.1)	6.1	(5.8 - 6.4)			+
Smoked cigars on one or more of the past 30 days	6.4	(4.8 - 8.0)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	2.3	(1.2 - 3.3)	3.5	(3.2 - 3.7)			+
Used any form of tobacco on one or more of the past 30 days	12.5	(10.1 - 14.8)	11.2	(10.8 - 11.7)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	64.1	(60.6 - 67.6)	65.7	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	49.5	(45.9 - 53.2)	44.1	(43.4 - 44.7)	+		
Smoking is allowed in the home	9.6	(7.6 - 11.6)	9.4	(9.0 - 9.8)			+
Taught about tobacco use in any classes this year	50.5	(46.1 - 54.9)	43.2	(42.3 - 44.1)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	58.2	(55.0 - 61.4)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	3.3	(2.0 - 4.5)	3.9	(3.7 - 4.1)			+
<u>Definitely did not think</u> that smokers have more friends	33.8	(30.9 - 36.7)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	71.9	(68.9 - 74.9)	72.9	(72.4 - 73.4)			+

Pinellas County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Pinellas County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	20.7	(18.0 - 23.4)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	11.6	(9.5 - 13.8)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	19.1	(13.4 - 24.8)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	7.2	(5.4 - 8.9)	11.1	(10.7 - 11.5)		+	
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	13.0	(10.7 - 15.4)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	67.8	(64.4 - 71.2)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	29.5	(26.5 - 32.5)	25.3	(24.8 - 25.8)	+		
Exercised to lose weight or to keep from gaining weight during the past 30 days	40.4	(37.2 - 43.7)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	28.4	(25.6 - 31.1)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	8.3	(6.3 - 10.3)	10.9	(10.4 - 11.3)		+	
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	20.3	(17.6 - 23.0)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	4.4	(3.1 - 5.8)	6.9	(6.6 - 7.2)		+	
Purposely hurt themselves without wanting to die, during the past 12 months	12.9	(10.3 - 15.4)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	22.7	(19.7 - 25.7)	22.0	(21.5 - 22.5)			+